

SNUTP 93-97, hep-th/9402049

BPS Dyon in a Weak Electromagnetic Field: Equations of Motion and Radiation Fields

Dongsu Bak

*Center for Theoretical Physics, Laboratory for Nuclear Science and
Department of Physics, Massachusetts Institute of Technology,
Cambridge, MA02139, USA.*

Choonkyu Lee

*Department of Physics and Center for Theoretical Physics,
Seoul National University, Seoul, 151-742, Korea*

Abstract

Dynamics of a BPS dyon in a weak, constant, electromagnetic field is studied through a perturbative analysis of appropriate non-linear field equations. The full Lorentz force law for a BPS dyon is established. Also derived are the radiation fields accompanying the motion.

The $SU(2)$ Yang-Mills-Higgs theory, spontaneously broken to $U(1)$ by an adjoint Higgs field, allows non-singular time-independent classical solutions of finite energy that carry nonzero magnetic charge [1, 2]. The same theory also admits dyons, solutions that carry both electric and magnetic charges [3]. In the so-called Prasad-Sommerfield limit [4] of vanishing scalar field potential, these monopoles or dyons become solutions of the first-order Bogomolny equations [5] or their suitable generalization [6]. A large number of papers have been written so far on these Bogomolny-Prasad-Sommerfield(BPS) monopoles, but most of them considered the *static* properties of the solutions only. A major exception here is the work by Manton and others [7], in which the scattering of slowly moving BPS monopoles is studied with the help of a geodesic approximation in the moduli space of static multi-monopole solutions. Very recently [8], we have also studied another important time-dependent phenomenon related to an accelerating BPS monopole, that of radiation made up of massless fields in the system. This involved a direct perturbative analysis of nonlinear field equations(also initiated by Manton [9]), and in the present work we report some additional results obtained by the same method.

The behavior of an electrically-neutral BPS monopole in a weak magnetic field $\mathbf{B}_0$ is known already from ref. [8]. We will here consider more general problem, i.e., the behavior of a BPS dyon (with magnetic charge g and electric charge q) in the presence of a weak, asymptotically uniform, electric and magnetic fields $(\mathbf{E}_0, \mathbf{B}_0)$. Following ref. [8], we will then have to construct an appropriate time-dependent solution to the nonlinear field equations, which describes the acceleration of the dyon and accompanying field deformations and radiation simultaneously. (For a more phenomenological approach to the related problem, see ref. [10]). To the first order in acceleration, we shall below set up a judicious perturbative scheme and then solve the necessary differential equations explicitly to exhibit the above-mentioned phenomena. A byproduct of our analysis is that the nonlinear field equations of the Yang-Mills-Higgs theory indeed require the center of a BPS dyon to move according to the naively expected equation of motion

$$\frac{d}{dt} \left(\frac{M_D \mathbf{v}}{\sqrt{1 - \mathbf{v}^2}} \right) = g(\mathbf{B}_0 - \mathbf{v} \times \mathbf{E}_0) + q(\mathbf{E}_0 + \mathbf{v} \times \mathbf{B}_0) \quad (1)$$

Note that our analysis is entirely in the classical field theory context, and so the electric charge q is not quantized. We set $c = 1$, and our metric convention is that with signature $(-+++)$.

We shall specify our system first. The Lagrangian density is ($a = 1, 2, 3$)

$$\mathcal{L} = -\frac{1}{4}G_a^{\mu\nu}G_{\mu\nu a} - \frac{1}{2}(D^\mu\phi)_a(D_\mu\phi)_a, \quad (2)$$

where

$$\begin{aligned} G_a^{\mu\nu} &= \partial^\mu A_a^\nu - \partial^\nu A_a^\mu + e\epsilon_{abc}A_b^\mu A_c^\nu, \\ (D_\mu\phi)^a &= \partial_\mu\phi^a + e\epsilon_{abc}A_{b\mu}\phi_c \equiv D_\mu^{ac}\phi_c, \end{aligned} \quad (3)$$

and the Higgs fields are subject to the asymptotic boundary condition $\phi^a\phi^a \rightarrow f^2 (\neq 0)$ as $r \rightarrow \infty$. The field equations read

$$D_\mu^{ac}G_c^{\mu\nu} = -e\epsilon_{abc}(D^\nu\phi)^b\phi^c, \quad (4a)$$

$$(D^\mu D_\mu\phi)^a = 0. \quad (4b)$$

If we here define the electric and magnetic charges by

$$q = \oint_{r=\infty} dS_i \frac{\phi^a}{|\phi|} E_i^a, \quad g = \oint_{r=\infty} dS_i \frac{\phi^a}{|\phi|} B_i^a \quad (5)$$

(with $E_i^a \equiv G_a^{0i}$, $B_i^a \equiv \frac{1}{2}\epsilon_{ijk}G_a^{jk}$), g must be of the form $\frac{4\pi}{e}n$ for some integer n by a topological reason while q can take on any continuous value. Remarkably, for given g and $q = g \tan \beta$, the static solutions to eqs.(4a) and (4b) with the lowest possible energy can be obtained by solving instead the first-order equations [5, 6]

$$B_i^a = \mp \cos \beta (D_i\phi)^a, \quad E_i^a = \mp \sin \beta (D_i\phi)^a, \quad (D_0\phi)^a = 0. \quad (6)$$

These are equations describing static BPS dyons and for $\beta = 0$ reduce to the Bogomolny equations for uncharged BPS monopoles:

$$B_i^a = \mp (D_i\phi)^a, \quad A_0^a = 0. \quad (7)$$

Actually all dyon solutions to eq.(6), denoted as $(\bar{\phi}^a(\mathbf{r}; \beta), \bar{A}_i^a(\mathbf{r}; \beta), \bar{A}_0^a(\mathbf{r}; \beta))$, can be obtained from the static monopole solutions $(\bar{\phi}^a(\mathbf{r}; \beta = 0), \bar{A}_i^a(\mathbf{r}; \beta = 0))$ satisfying eq.(7). This is achieved by the simple substitution [11]

$$\bar{\phi}^a(\mathbf{r}; \beta) = [\bar{\phi}^a(\mathbf{y}; \beta = 0)]_{\mathbf{y}=\mathbf{r} \cos \beta}, \quad (8a)$$

$$\bar{A}_i^a(\mathbf{r}; \beta) = \cos \beta [\bar{A}_i^a(\mathbf{y}; \beta = 0)]_{\mathbf{y}=\mathbf{r} \cos \beta}, \quad (8b)$$

$$\begin{aligned} \bar{A}_0^a(\mathbf{r}; \beta) &= \mp \sin \beta [\bar{\phi}^a(\mathbf{y}; \beta = 0)]_{\mathbf{y}=\mathbf{r} \cos \beta} \\ &= \mp \sin \beta \bar{\phi}^a(\mathbf{r}; \beta). \end{aligned} \quad (8c)$$

Especially, by making this substitution with the well-known BPS one-monopole solution (with $g = \mp \frac{4\pi}{e}$ and mass $M = \frac{4\pi}{e}f$)

$$\begin{aligned}\bar{\phi}^a(\mathbf{r}; \beta = 0) &= \pm \hat{\mathbf{r}}_a f \left[\coth m_v r - \frac{1}{m_v r} \right] , \quad (m_v = ef) \\ \bar{A}_i^a(\mathbf{r}; \beta = 0) &= \epsilon_{ail} \frac{\hat{\mathbf{r}}_l}{er} \left[1 - \frac{m_v r}{\sinh m_v r} \right] ,\end{aligned}\tag{9}$$

we immediately obtain the spherically symmetric BPS dyon solution [4] with $g = \mp \frac{4\pi}{e}$, $q = \mp \frac{4\pi}{e} \tan \beta$ and mass $M_D = \frac{4\pi}{e} \frac{f}{\cos \beta}$. In this paper we will focus on time-dependent dynamics of this dyon when there is a weak uniform electromagnetic field asymptotically, viz., under the condition that

$$r \rightarrow \infty : \quad \frac{\phi^a}{|\phi|} B_i^a \rightarrow (\mathbf{B}_0)_i , \quad \frac{\phi^a}{|\phi|} E_i^a \rightarrow (\mathbf{E}_0)_i ,\tag{10}$$

It is impractical to look for exact solutions to the full field equations (4a) and (4b) appropriate to this circumstance. But, to the first order in $\mathbf{B}_0$ and $\mathbf{E}_0$ (and hence to the first order in the dyon acceleration $\mathbf{a}$ also), the corresponding approximate solutions to eqs.(4a) and (4b) can be constructed explicitly.

Let us first generalize the result of ref. [8] for a BPS monopole (with $q = 0$) by allowing the asymptotic electric field $\mathbf{E}_0$ as well. Assuming that the monopole has zero velocity at $t = 0$, we may then write the ansatz

$$\begin{aligned}\phi^a(\mathbf{r}, t) &= \bar{\phi}^a(\mathbf{r}'; \beta = 0) + \Pi^a(\mathbf{r}') \quad (\equiv \tilde{\phi}^a(\mathbf{r}'; \beta = 0)) , \\ \mathbf{A}_i^a(\mathbf{r}, t) &= \bar{A}_i^a(\mathbf{r}'; \beta = 0) + \alpha_i^a(\mathbf{r}') \quad (\equiv \tilde{A}_i^a(\mathbf{r}'; \beta = 0)) , \\ \mathbf{A}_0^a(\mathbf{r}, t) &= -ta_i \bar{A}_i^a(\mathbf{r}'; \beta = 0) + \alpha_0^a(\mathbf{r}')\end{aligned}\tag{11}$$

with $\mathbf{r}' \equiv \mathbf{r} - \frac{1}{2}\mathbf{a}t^2$, $\mathbf{a}$ being the expected acceleration of the monopole. Here, $(\bar{\phi}^a, \bar{A}_i^a)$ denotes the BPS one-monopole solutions in eq.(9) and the presence of the term $\alpha_0^a(\mathbf{r}')$ distinguishes the present form from that of ref. [8]. Our approximation is to neglect any terms beyond $O(a)$ (here $a \equiv |\mathbf{B}_0|$, $|\mathbf{E}_0|$ or $|\mathbf{a}|$) in Π^a , α_i^a and α_0^a , and in this approximation the consistency of our assumption that these functions depend on time only through $\mathbf{r}'$ can be established. Then, from the field equations (4a) and (4b), we can deduce the following set of equations:

$$(\bar{D}^i \alpha^j)^a - (\bar{D}^j \alpha^i)^a = \mp \epsilon^{ijk} \left(\bar{D}_k^{ab} \Pi^b + e \epsilon_{abc} \alpha_k^b \bar{\phi}^c + a_k \bar{\phi}^a \right) ,\tag{12a}$$

$$(\bar{D}_i \bar{D}_i \alpha_0)^a = -e^2 \epsilon_{abc} \epsilon_{bdf} \bar{\phi}_c \bar{\phi}_f \alpha_0^d ,\tag{12b}$$

where we have defined $\bar{D}_i^{ac} = \partial_i \delta_{ac} + e \epsilon_{abc} \bar{A}_i^b$. Since $\bar{B}_i^a \equiv \frac{1}{2} \epsilon_{ijk} \bar{G}_a^{jk} = \mp (\bar{D}_i \bar{\phi})^a$, eq.(12a) is equivalent to [9]

$$B_i^a = \mp (D_i + a_i)^{ab} \phi_b , \quad (13)$$

while the field strength E_i^a can be expressed as

$$E_i^a = -t a_j \bar{G}_a^{ij} + (\bar{D}_i \alpha_0)^a . \quad (14)$$

Then, comparing these with our asymptotic requirement (10), we can immediately conclude that

- (i) $\mathbf{B}_0 = \mp \mathbf{a} f$ or, equivalently, $M \mathbf{a} = g \mathbf{B}_0$ ($M = \frac{4\pi}{e} f$ is the monopole mass), and
- (ii) $\alpha_0^a(\mathbf{r}') \rightarrow \pm \mathbf{E}_0 \cdot \mathbf{r}' \hat{\mathbf{r}}'_a$ as $r' = |\mathbf{r}'| \rightarrow \infty$.

The functions $\Pi^a(\mathbf{r}')$ and $\alpha_i^a(\mathbf{r}')$, which are solutions to eq.(12a), have been found already in ref. [8]. Using those results, we can equate $\phi^a(\mathbf{r}, t)$ and $A_i^a(\mathbf{r}, t)$ to the first order in $\mathbf{B}_0$ and $\mathbf{E}_0$ to the expressions (here $\mathbf{a} = \mp \mathbf{B}_0/f$)

$$\tilde{\phi}^a(\mathbf{r}'; \beta = 0) = \pm \hat{\mathbf{r}}'_a \left[f \left(\coth m_v r' - \frac{1}{m_v r'} \right) + \frac{1}{2e} \hat{\mathbf{r}}' \cdot \mathbf{a} \left(1 - \frac{m_v r'}{\sinh m_v r'} \right) \right] \pm \frac{1}{2} a_a \frac{f r'}{\sinh m_v r'} , \quad (15a)$$

$$\tilde{A}_i^a(\mathbf{r}'; \beta = 0) = \epsilon_{aij} \frac{\hat{\mathbf{r}}'_j}{e r'} \left(1 - \frac{m_v r'}{\sinh m_v r'} \right) + \frac{f r'}{2} \left(\coth m_v r' - \frac{1}{m_v r'} \right) \epsilon_{ijk} \hat{\mathbf{r}}'_j a_k \hat{\mathbf{r}}'_a , \quad (15b)$$

respectively. Remarkably the closed-form solution to eq.(12b) which satisfies the above asymptotic boundary condition can also be found:

$$\alpha_0^a(\mathbf{r}') = \pm (\coth m_v r') \mathbf{E}_0 \cdot \mathbf{r}' \hat{\mathbf{r}}'_a \pm \frac{r'}{\sinh m_v r'} [(\mathbf{E}_0)_a - (\mathbf{E}_0 \cdot \hat{\mathbf{r}}') \hat{\mathbf{r}}'_a] . \quad (16)$$

The functions $\alpha_0^a(\mathbf{r}')$ here describe necessary field deformations in the presence of an asymptotic electric field $\mathbf{E}_0$; but, in association with this term, there is no additional electromagnetic or Higgs scalar radiation (beyond those already present in association with the asymptotic magnetic field $\mathbf{B}_0$). On the other hand, this solution in the presence of nonzero $\mathbf{B}_0$ and $\mathbf{E}_0$ will be important to discuss the Lorentz boost property. See below.

If one considers a Lorentz boost transformation of the solution specified by eqs.(11), (15a), (15b) and (16), viz.,

$$\begin{aligned}
\phi^{*a}(\mathbf{r}, t) &= \phi^a(\mathbf{r}^*, t^*) , \\
A_i^{*a}(\mathbf{r}, t) &= A_i^a(\mathbf{r}^*, t^*) + \left(\frac{1}{\sqrt{1 - \mathbf{v}_0^2}} - 1 \right) (\hat{\mathbf{v}}_0^j A_j^a(\mathbf{r}^*, t^*)) \hat{\mathbf{v}}_0^i - \frac{1}{\sqrt{1 - \mathbf{v}_0^2}} v_0^i A_0^a(\mathbf{r}^*, t^*) , \\
A_0^{*a}(\mathbf{r}, t) &= \frac{1}{\sqrt{1 - \mathbf{v}_0^2}} \left\{ A_0^a(\mathbf{r}^*, t^*) - v_0^j A_j^a(\mathbf{r}^*, t^*) \right\} , \\
\left(\mathbf{r}^* = \mathbf{r} + \left(\frac{1}{\sqrt{1 - \mathbf{v}_0^2}} - 1 \right) (\hat{\mathbf{v}}_0 \cdot \mathbf{r}) \hat{\mathbf{v}}_0 - \frac{1}{\sqrt{1 - \mathbf{v}_0^2}} \mathbf{v}_0 t ; \quad t^* = \frac{t - \mathbf{v}_0 \cdot \mathbf{r}}{\sqrt{1 - \mathbf{v}_0^2}} \right)
\end{aligned} \tag{17}$$

the result will be another solution to the field equations. It describes an accelerating BPS monopole with finite initial velocity $\mathbf{v}_0$ in the presence of a uniform electromagnetic field $(\mathbf{E}_0^*, \mathbf{B}_0^*)$, where

$$\begin{aligned}
\mathbf{E}_0^* &= (\hat{\mathbf{v}}_0 \cdot \mathbf{E}_0) \hat{\mathbf{v}}_0 + \frac{1}{\sqrt{1 - \mathbf{v}_0^2}} [\mathbf{E}_0 - (\hat{\mathbf{v}}_0 \cdot \mathbf{E}_0) \hat{\mathbf{v}}_0 - \mathbf{v}_0 \times \mathbf{B}_0] , \\
\mathbf{B}_0^* &= (\hat{\mathbf{v}}_0 \cdot \mathbf{B}_0) \hat{\mathbf{v}}_0 + \frac{1}{\sqrt{1 - \mathbf{v}_0^2}} [\mathbf{B}_0 - (\hat{\mathbf{v}}_0 \cdot \mathbf{B}_0) \hat{\mathbf{v}}_0 + \mathbf{v}_0 \times \mathbf{E}_0] .
\end{aligned} \tag{18}$$

[Especially, by choosing $\mathbf{B}_0 = \mathbf{v}_0 \times \mathbf{E}_0$, one can obtain the solution appropriate to a BPS monopole with a finite initial velocity $\mathbf{v}_0$ in the presence of some nonzero electric field $\mathbf{E}_0^*$ (but $\mathbf{B}_0^* = 0$).] The trajectory assumed by the center of this monopole will be determined by the condition $\mathbf{r}^* - \frac{1}{2} \left(\frac{g\mathbf{B}_0}{M} \right) (t^*)^2 = 0$, or by

$$\begin{aligned}
\frac{d^2 \mathbf{r}^*}{dt^{*2}} &= \frac{g}{M} \mathbf{B}_0 \\
&= (\hat{\mathbf{v}}_0 \cdot \mathbf{B}^*) \hat{\mathbf{v}}_0 + \frac{1}{\sqrt{1 - \mathbf{v}_0^2}} [\mathbf{B}_0^* - (\hat{\mathbf{v}}_0 \cdot \mathbf{B}_0^*) \hat{\mathbf{v}}_0 - \mathbf{v}_0 \times \mathbf{E}_0^*] ,
\end{aligned} \tag{19}$$

where we have used eq.(18). At the same time, we have

$$\frac{d^2 \mathbf{r}^*}{dt^{*2}} = \frac{(\hat{\mathbf{v}}_0 \cdot \frac{d\mathbf{v}}{dt}) \hat{\mathbf{v}}_0}{(1 - \mathbf{v}_0^2)^{3/2}} + \frac{1}{(1 - \mathbf{v}_0^2)} \left\{ \frac{d\mathbf{v}}{dt} - \left(\hat{\mathbf{v}}_0 \cdot \frac{d\mathbf{v}}{dt} \right) \hat{\mathbf{v}}_0 \right\} + O(a^2) , \tag{20}$$

(here, $\mathbf{v}(t) = \frac{d\mathbf{r}(t)}{dt}$ ($= \mathbf{v}_0 + O(a)$) denotes the velocity of the monopole), and consequently eq.(19) can be recast as

$$\frac{d}{dt} \left(\frac{M\mathbf{v}}{\sqrt{1 - \mathbf{v}^2}} \right) = g [\mathbf{B}_0^* - \mathbf{v} \times \mathbf{E}^*] + O(a^2) . \tag{21}$$

Eq.(1) is thus confirmed for an uncharged monopole.

A similar analysis can also be carried out for a BPS dyon, at least to the first order in the asymptotic electromagnetic fields $\mathbf{E}_0$ and $\mathbf{B}_0$ (and hence in the dyon acceleration $\mathbf{a}$). Again choosing the reference frame in such a way that the dyon may have zero velocity at $t = 0$, we may generalize the ansatz (11) as

$$\begin{aligned}\phi^a(\mathbf{r}, t) &= \tilde{\phi}^a(\mathbf{r}'; \beta) , \\ A_i^a(\mathbf{r}, t) &= -ta_i \bar{A}_0^a(\mathbf{r}'; \beta) + \tilde{A}_i^a(\mathbf{r}'; \beta) , \\ A_0^a(\mathbf{r}, t) &= -ta_i \bar{A}_i^a(\mathbf{r}'; \beta) + \tilde{A}_0^a(\mathbf{r}'; \beta)\end{aligned}\tag{22}$$

with

$$\begin{aligned}\tilde{\phi}^a(\mathbf{r}'; \beta) &= \bar{\phi}^a(\mathbf{r}'; \beta) + \Pi^a(\mathbf{r}'; \beta) , \\ \tilde{A}_i^a(\mathbf{r}'; \beta) &= \bar{A}_i^a(\mathbf{r}'; \beta) + \alpha_i^a(\mathbf{r}'; \beta) , \\ \tilde{A}_0^a(\mathbf{r}'; \beta) &= \mp \sin \beta \bar{\phi}^a(\mathbf{r}'; \beta) + \alpha_0^a(\mathbf{r}'; \beta) ,\end{aligned}\tag{23}$$

where $\mathbf{r}' = \mathbf{r} - \frac{1}{2}\mathbf{a}t^2$. Here the functions $\bar{\phi}^a(\mathbf{r}; \beta)$, $\bar{A}_i^a(\mathbf{r}; \beta)$ and $\bar{A}_0^a(\mathbf{r}; \beta)$ represent the static dyon solution in eqs.(8a)-(8c), and the yet-to-be-determined functions $(\Pi^a, \alpha_i^a, \alpha_0^a)$ are assumed to be $O(a)$. Notice the specific way the time coordinate t enters our ansatz. The appearance of the piece $-ta_i \bar{A}_0^a(\mathbf{r}'; \beta)$ or $-ta_i \bar{A}_i^a(\mathbf{r}'; \beta)$ in eq.(22) may well be understood from the viewpoint of the instantaneous Lorentz boost. Also see the relationship (8c) for a possible motivation for writing $\tilde{A}_0^a(\mathbf{r}'; \beta)$ as in eq.(23).

The consistency of the above ansatz can be verified explicitly; inserting the ansatz (22) into the full field equations leads to differential equations whose sole dependent variables are $\mathbf{r}'$. Explicitly, we obtain from eq.(4a)

$$(\tilde{D}_j + a_j)^{ac} \tilde{G}_c^{ji} + e\epsilon_{abc} \tilde{A}_0^b (\tilde{D}_i \tilde{A}_0)^c = -e\epsilon_{abc} (\tilde{D}_i \tilde{\phi})^b \tilde{\phi}^c ,\tag{24a}$$

$$\tilde{D}_i^{ab} (\tilde{D}_i \tilde{A}_0 + a_i \tilde{A}_0)^b + e^2 \epsilon_{abc} \epsilon_{bdf} \tilde{\phi}^c \tilde{A}_0^d \tilde{\phi}^f = 0 ,\tag{24b}$$

and from eq.(4b)

$$\tilde{D}_i^{ab} (\tilde{D}_i \tilde{\phi} + a_i \tilde{\phi})^b + e^2 \epsilon_{abc} \epsilon_{bdf} \tilde{A}_0^c \tilde{A}_0^d \tilde{\phi}^f = 0 ,\tag{24c}$$

where $\tilde{D}_i^{ab} = [D_i^{ab}]_{A_i^a \rightarrow \tilde{A}_i^a}$, $\tilde{G}_c^{ji} = [G_c^{ji}]_{A_i^a \rightarrow \tilde{A}_i^a}$, and the suppressed dependent variable is $\mathbf{r}'$. Then, identifying $\tilde{A}_0^a(\mathbf{r}'; \beta)$ as in eq.(23), we found that these three equations are satisfied (to $O(a)$) as long as the following conditions are fulfilled:

$$\tilde{B}_i^a = \mp (\tilde{D}_i + a_i)^{ab} (\cos \beta \tilde{\phi}^b \pm \tan \beta \alpha_0^b) ,\tag{25a}$$

$$(\tilde{D}_i \tilde{D}_i \alpha_0)^a = -e^2 \cos^2 \beta \epsilon_{abc} \epsilon_{bdf} \bar{\phi}^c \bar{\phi}^f \alpha_0^d .\tag{25b}$$

Notice that, for $\beta = 0$, these reduce to the corresponding equations for an uncharged BPS monopole, i.e., eqs.(12b) and (13). In the present case the field strength E_i^a to $O(a)$ becomes

$$E_i^a(\mathbf{r}, t) = -ta_j \bar{G}_a^{ij} + (\tilde{D}_i + a_i)^{ab} \tilde{A}_0^b. \quad (26)$$

The acceleration $\mathbf{a}$ above is not arbitrary but fixed in terms of the given asymptotic electromagnetic fields. For this purpose the asymptotic condition (10) may be used with the field strengths given by eqs.(25a) and (26). Actually we will also assume that $\frac{\phi^a}{|\phi|}(\tilde{D}_i \tilde{\phi})^a \rightarrow 0$ as $r' \rightarrow \infty$; this can be viewed as our additional asymptotic condition. Moreover, according to our lesson from the uncharged monopole case, it is quite natural to suppose that $\alpha_0^a(\mathbf{r}'; \beta) \rightarrow \cos \beta \boldsymbol{\mathcal{E}}_0 \cdot \mathbf{r}' \hat{\mathbf{r}}'_a$ (for some constant vector $\boldsymbol{\mathcal{E}}_0$) as $r' \rightarrow \infty$. Now, from eq.(25a) and the asymptotic conditions,

$$\mathbf{B}_0 = \mp f \cos \beta \mathbf{a} \mp \sin \beta \boldsymbol{\mathcal{E}}_0, \quad (27)$$

while eq.(26) and the asymptotic conditions lead to

$$\mathbf{E}_0 = \mp f \sin \beta \mathbf{a} \pm \cos \beta \boldsymbol{\mathcal{E}}_0. \quad (28)$$

We thus have

$$\boldsymbol{\mathcal{E}}_0 = \mp (\sin \beta \mathbf{B}_0 - \cos \beta \mathbf{E}_0), \quad \mathbf{a} = \mp \frac{1}{f} (\cos \beta \mathbf{B}_0 + \sin \beta \mathbf{E}_0), \quad (29)$$

and especially, recalling that the dyon has mass $M_D = \frac{4\pi}{e} \frac{f}{\cos \beta}$, $g = \mp \frac{4\pi}{e}$ and $q = \mp \frac{4\pi}{e} \tan \beta$, the second of these can be recast as

$$M_D \mathbf{a} = g \mathbf{B}_0 + q \mathbf{E}_0. \quad (30)$$

This corresponds to the formula (1) in dyon's instantaneous rest frame. The equation of motion in the general form (1) can be secured if the Lorentz boost transformation (17) is considered for the above solution. This proceeds in much the same way as in eqs.(17)-(21), and so we shall not repeat it here.

We will now find the explicit solution to eqs.(25a) and (25b). Here it is convenient to introduce rescaled quantities

$$\mathbf{y} = \mathbf{r}' \cos \beta, \quad \mathcal{A}_i^a(\mathbf{y}) = \frac{1}{\cos \beta} \tilde{A}_i^a\left(\mathbf{r}' = \frac{\mathbf{y}}{\cos \beta}; \beta\right). \quad (31)$$

Then, with the help of the relationships (8a)-(8c), we can rewrite eqs.(25a) and (25b) as

$$\mathcal{B}_i^a = \mp \left(\mathcal{D}_i^{(y)} + \frac{a_i}{\cos \beta} \right)^{ab} \left(\tilde{\phi}^b \pm \frac{\sin \beta}{\cos^2 \beta} \alpha_0^b \right), \quad (32a)$$

$$(\bar{D}_i^{(y)} \bar{D}_i^{(y)} \alpha_0)^a = -e^2 \epsilon_{abc} \epsilon_{bdf} \bar{\phi}^c(\mathbf{y}; \beta = 0) \bar{\phi}^f(\mathbf{y}; \beta = 0) \alpha_0^d, \quad (32b)$$

where $\bar{D}_i^{(y)ac} \equiv \frac{\partial}{\partial y_i} \delta_{ac} + e \epsilon_{abc} \bar{A}_i^b(\mathbf{y}; \beta = 0)$, $\mathcal{D}_i^{(y)ac} \equiv \frac{\partial}{\partial y_i} \delta_{ac} + e \epsilon_{abc} \mathcal{A}_i^b(\mathbf{y})$, and $\mathcal{B}_i^a(\mathbf{y})$ denotes the magnetic field strength obtained from the vector potential $\mathcal{A}_i^a(\mathbf{y})$. Eq.(32b) coincides with our earlier equation(eq.(12b)), and so the solution with the desired asymptotic behavior is immediately identified with (here $y = |\mathbf{y}| = r' \cos \beta$)

$$\alpha_0^a = (\coth m_v y) \mathcal{E}_0 \cdot \mathbf{y} \hat{\mathbf{y}}_a + \frac{y}{\sinh m_v y} [(\mathcal{E}_0)_a - (\mathcal{E}_0 \cdot \hat{\mathbf{y}}) \hat{\mathbf{y}}_a]. \quad (33)$$

Moreover, comparing eq.(32a) with eq.(13), we are led to write

$$\begin{aligned} \mathcal{A}_i^a &= [\tilde{A}_i^a(\mathbf{y}; \beta = 0)]_{\mathbf{a} \rightarrow \frac{\mathbf{a}}{\cos \beta}} + \hat{\alpha}_i^a(\mathbf{y}), \\ \tilde{\phi}^a &= \mp \frac{\sin \beta}{\cos^2 \beta} \alpha_0^a + [\tilde{\phi}^a(\mathbf{y}; \beta = 0)]_{\mathbf{a} \rightarrow \frac{\mathbf{a}}{\cos \beta}} + \hat{\Pi}^a(\mathbf{y}), \end{aligned} \quad (34)$$

where $\hat{\alpha}_i^a(\mathbf{y})$ and $\hat{\Pi}^a(\mathbf{y})$ now satisfy the homogeneous equation (cf. eq.(12a))

$$(\bar{D}^{(y)i} \hat{\alpha}^j)^a - (\bar{D}^{(y)j} \hat{\alpha}^i)^a = \mp \epsilon^{ijk} (\bar{D}_k^{(y)ab} \hat{\Pi}^b + e \epsilon_{abc} \hat{\alpha}_k^b \bar{\phi}^c(\mathbf{y}; \beta = 0)), \quad (35)$$

the functions $\tilde{\phi}^a(\mathbf{y}; \beta = 0)$ and $\tilde{A}_i^a(\mathbf{y}; \beta = 0)$ are given explicitly in eqs(15a) and (15b), and α_0^a in eq.(33). Since the function α_0^a increases linearly for large r' , we must here include a suitable homogeneous solution ($\hat{\alpha}_i^a, \hat{\Pi}^a$) so that our asymptotic condition $\lim_{r' \rightarrow \infty} \frac{\phi^a}{|\phi|} (\tilde{D}_i \tilde{\phi})^a = 0$ may be fulfilled.

The general, normalizable or nonnormalizable, solutions of eq. (35) have been known for some time [12]. Because of the above-mentioned asymptotic condition, we may here demand $\hat{\Pi}^a(\mathbf{y})$ to behave such as

$$y \rightarrow \infty : \quad \hat{\Pi}^a(\mathbf{y}) \rightarrow \pm \frac{\sin \beta}{\cos^2 \beta} \mathcal{E}_0 \cdot \mathbf{y} \hat{\mathbf{y}}_a. \quad (36)$$

Then, using the result of ref. [12], we find that the appropriate solution to eq. (35) is given by

$$\begin{aligned}\hat{\alpha}_i^a(\mathbf{y}) &= \epsilon_{ail} \frac{\partial}{\partial y^l} \left(\frac{y}{\sinh m_v y} V \right) + (1 - \cosh m_v y) \hat{\mathbf{y}}_a \epsilon_{ilm} \hat{\mathbf{y}}_l \frac{\partial}{\partial y^m} \left(\frac{y}{\sinh m_v y} V \right), \\ \hat{\Pi}^a(\mathbf{y}) &= \mp \hat{\mathbf{y}}^a \frac{\partial}{\partial y} (y \coth m_v y V) \mp \left(\frac{\partial}{\partial y^a} - \hat{\mathbf{y}}_a \frac{\partial}{\partial y} \right) \left(\frac{y}{\sinh m_v y} V \right)\end{aligned}\quad (37)$$

with $V = -\frac{\sin \beta}{2 \cos^2 \beta} \boldsymbol{\mathcal{E}}_0 \cdot \mathbf{y}$. Based on these findings, the functions $\tilde{\phi}^a(\mathbf{r}'; \beta)$, $\tilde{A}_i^a(\mathbf{r}'; \beta)$ and $\tilde{A}_0^a(\mathbf{r}'; \beta)$ in eq. (22) are explicitly given as

$$\begin{aligned}\tilde{\phi}^a(\mathbf{r}'; \beta) &= \pm \hat{\mathbf{r}}'_a \left[f \left\{ \coth[(m_v \cos \beta) r'] - \frac{1}{(m_v \cos \beta) r'} \right\} \right. \\ &\quad \left. + \frac{\hat{\mathbf{r}}' \cdot \mathbf{a}}{2e \cos \beta} \left\{ 1 - \frac{(m_v \cos \beta) r'}{\sinh[(m_v \cos \beta) r']} \right\} \right] \\ &\quad \pm \frac{f}{2} a_a \frac{r'}{\sinh[(m_v \cos \beta) r']} \mp \frac{1}{2} (\boldsymbol{\mathcal{E}}_0 \cdot \mathbf{r}') \hat{\mathbf{r}}'_a \frac{(m_v \sin \beta) r'}{\sinh^2[(m_v \cos \beta) r']} \\ &\quad \mp \frac{\tan \beta}{2} [(\boldsymbol{\mathcal{E}}_0)_a - (\boldsymbol{\mathcal{E}}_0 \cdot \hat{\mathbf{r}}') \hat{\mathbf{r}}'_a] \frac{r'}{\sinh[(m_v \cos \beta) r']}, \\ \tilde{A}_i^a(\mathbf{r}'; \beta) &= \epsilon_{aij} \frac{\hat{\mathbf{r}}'_j}{e r'} \left\{ 1 - \frac{(m_v \cos \beta) r'}{\sinh[(m_v \cos \beta) r']} \right\} \\ &\quad + \frac{f \cos \beta}{2} r' \left\{ \coth[(m_v \cos \beta) r'] - \frac{1}{(m_v \cos \beta) r'} \right\} \epsilon^{ijk} \hat{\mathbf{r}}'_j a_k \hat{\mathbf{r}}'_a \\ &\quad - \frac{\sin \beta}{2} \epsilon_{ail} \frac{\partial}{\partial r'^l} \left\{ \frac{r' (\boldsymbol{\mathcal{E}}_0 \cdot \mathbf{r}')}{\sinh[(m_v \cos \beta) r']} \right\} \\ &\quad - \frac{\sin \beta}{2} \frac{r' \{1 - \cosh[(m_v \cos \beta) r']\}}{\sinh[(m_v \cos \beta) r']} \hat{\mathbf{r}}'_a \epsilon_{ilm} \hat{\mathbf{r}}'_l (\boldsymbol{\mathcal{E}}_0)_m, \\ \tilde{A}_0^a(\mathbf{r}'; \beta) &= \mp \sin \beta \tilde{\phi}^a(\mathbf{r}'; \beta) + \cos \beta \coth[(m_v \cos \beta) r'] \boldsymbol{\mathcal{E}}_0 \cdot \mathbf{r}' \hat{\mathbf{r}}'_a \\ &\quad + \frac{(\cos \beta) r'}{\sinh[(m_v \cos \beta) r']} [(\boldsymbol{\mathcal{E}}_0)_a - (\boldsymbol{\mathcal{E}}_0 \cdot \hat{\mathbf{r}}') \hat{\mathbf{r}}'_a]\end{aligned}\quad (38)$$

with the constant vectors $\boldsymbol{\mathcal{E}}_0$ and $\mathbf{a}$ determined by eq. (29). Notice that these functions are everywhere regular.

In eqs. (22) and (38) we have the explicit solution appropriate to an accelerating BPS dyon. Its asymptotic behavior is of particular interest since it carries information on emitted radiation. Let us concentrate on the asymptotic behaviors of the corresponding gauge-covariant quantities. A short calculation using eqs. (22) and (38), while ignoring exponentially small terms, produces the following results (valid for $(m_v \cos \beta)r' \gg 1$):

$$G_a^{ij}(\mathbf{r}, t) \sim \epsilon^{ijk} \left\{ -\frac{1}{er'^2} \hat{\mathbf{r}}'_k + \frac{1}{2er'} [a_k + \hat{\mathbf{r}}'_k (\hat{\mathbf{r}}' \cdot \mathbf{a})] + \frac{\tan \beta}{e} \frac{t(\hat{\mathbf{r}}' \times \mathbf{a})_k}{r'^2} - f \cos \beta a_k - \sin \beta (\mathcal{E}_0)_k \right\} \hat{\mathbf{r}}'_a, \quad (39a)$$

$$G_a^{0i}(\mathbf{r}, t) \sim \left\{ -\frac{\tan \beta}{e} \frac{\hat{\mathbf{r}}'_i}{r'^2} + \frac{\tan \beta}{2e} [a_i + \hat{\mathbf{r}}'_i (\hat{\mathbf{r}}' \cdot \mathbf{a})] - \frac{t(\hat{\mathbf{r}}' \times \mathbf{a})_i}{er'^2} - f \sin \beta a_i + \cos \beta (\mathcal{E}_0)_i \right\} \hat{\mathbf{r}}'_a, \quad (39b)$$

$$(D^i \phi)^a(\mathbf{r}, t) \sim \pm \frac{1}{e \cos \beta} \left\{ \frac{\hat{\mathbf{r}}'_i}{r'^2} + \frac{a_i - \hat{\mathbf{r}}'_i (\hat{\mathbf{r}}' \cdot \mathbf{a})}{2r'} \right\} \hat{\mathbf{r}}'_a, \quad (39c)$$

$$(D^0 \phi)^a(\mathbf{r}, t) \sim \pm \frac{1}{e \cos \beta} \frac{t \mathbf{a} \cdot \hat{\mathbf{r}}'}{r'^2} \hat{\mathbf{r}}'_a. \quad (39d)$$

The asymptotic electromagnetic fields also follow from these expressions by setting $G_a^{ij}(\mathbf{r}, t) \sim \epsilon^{ijk} B_k^{em}(\mathbf{r}, t) \hat{\mathbf{r}}'_a$ and $G_a^{0i}(\mathbf{r}, t) \sim E_i^{em}(\mathbf{r}, t) \hat{\mathbf{r}}'_a$. Non-trivial expressions we have in eqs. (39c) and (39d) can be interpreted as the long-range effects associated with the massless Higgs scalar in this theory.

Now, as in Ref. [8], it will be convenient to rewrite the above expressions using the “retarded” distance $\mathbf{R} = \mathbf{r} - \frac{1}{2} \mathbf{a} t_{ret}^2$, where t_{ret} is determined (for given $\mathbf{r}$ and t) through the implicit equation $t - t_{ret} = |\mathbf{r} - \frac{1}{2} \mathbf{a} t_{ret}^2| \equiv R$. Here, noting that $\frac{1}{2} |\mathbf{a}| t_{ret}^2 \ll R$ for sufficiently small acceleration, we have

$$\mathbf{r}' = \mathbf{R} - \mathbf{v}_{ret} R - \frac{1}{2} \mathbf{a} R^2, \quad (\mathbf{v}_{ret} \equiv \mathbf{a} t_{ret}) \quad (40)$$

and $r' = R(1 - \hat{\mathbf{R}} \cdot \mathbf{v}_{ret} - \frac{1}{2} R \hat{\mathbf{R}} \cdot \mathbf{a})$. With these and the identifications $g = \mp \frac{4\pi}{e}$ and

$q = \mp \frac{4\pi}{e} \tan \beta$, it is now a simple matter to recast the above asymptotic fields as

$$\begin{aligned} \mathbf{B}^{em}(\mathbf{r}, t) \sim & \frac{g}{4\pi} \frac{\hat{\mathbf{R}} - \mathbf{v}_{ret}}{(1 - \hat{\mathbf{R}} \cdot \mathbf{v}_{ret})^3 R^2} + \frac{g}{4\pi} \frac{\hat{\mathbf{R}} \times (\hat{\mathbf{R}} \times \mathbf{a})}{R} \\ & - \frac{q}{4\pi} \frac{\hat{\mathbf{R}} \times \mathbf{v}_{ret}}{R^2} - \frac{q}{4\pi} \frac{\hat{\mathbf{R}} \times \mathbf{a}}{R} + \mathbf{B}_0, \end{aligned} \quad (41a)$$

$$\begin{aligned} \mathbf{E}^{em}(\mathbf{r}, t) \sim & \frac{q}{4\pi} \frac{\hat{\mathbf{R}} - \mathbf{v}_{ret}}{(1 - \hat{\mathbf{R}} \cdot \mathbf{v}_{ret})^3 R^2} + \frac{q}{4\pi} \frac{\hat{\mathbf{R}} \times (\hat{\mathbf{R}} \times \mathbf{a})}{R} \\ & + \frac{g}{4\pi} \frac{\hat{\mathbf{R}} \times \mathbf{v}_{ret}}{R^2} + \frac{g}{4\pi} \frac{\hat{\mathbf{R}} \times \mathbf{a}}{R} + \mathbf{E}_0, \end{aligned} \quad (41b)$$

$$(D^i \phi)^a(\mathbf{r}, t) \sim \left\{ \frac{g_s}{4\pi} \frac{\hat{\mathbf{R}}_i - v_{ret,i}}{(1 - \hat{\mathbf{R}} \cdot \mathbf{v}_{ret})^3 R^2} + \frac{g_s}{4\pi} \frac{(\hat{\mathbf{R}} \cdot \mathbf{a}) \hat{\mathbf{R}}_i}{R} \right\} \hat{\mathbf{r}}'_a, \quad (41c)$$

$$(D^0 \phi)^a(\mathbf{r}, t) \sim \left\{ \frac{g_s}{4\pi} \frac{\hat{\mathbf{R}} \cdot \mathbf{v}_{ret}}{R^2} + \frac{g_s}{4\pi} \frac{\hat{\mathbf{R}} \cdot \mathbf{a}}{R} \right\} \hat{\mathbf{r}}'_a, \quad (41d)$$

where we have defined $g_s = \pm \frac{4\pi}{e \cos \beta}$, and $\mathbf{E}_0$ and $\mathbf{B}_0$ are the asymptotic uniform electromagnetic fields given through eqs. (27) and (28). In these expressions the $O(R^{-2})$ terms make the near-zone fields associated with electromagnetic or Higgs scalar force. Observe that the near-zone fields we have obtained for $\mathbf{B}^{em}$ and $\mathbf{E}^{em}$ agree with the naively expected results on the basis of the classical Maxwell theory and the duality argument [13]. The same can be said also for the radiation fields, which we discuss below.

The $O(R^{-1})$ terms in eqs.(41a)-(41d) describe the radiation fields accompanying an accelerating BPS dyon. Explicitly we have

$$\begin{aligned} \mathbf{B}_{rad} &= \frac{g}{4\pi} \frac{\hat{\mathbf{R}} \times (\hat{\mathbf{R}} \times \mathbf{a})}{R} + \frac{q}{4\pi} \frac{\mathbf{a} \times \hat{\mathbf{R}}}{R}, \\ \mathbf{E}_{rad} &= \frac{q}{4\pi} \frac{\hat{\mathbf{R}} \times (\hat{\mathbf{R}} \times \mathbf{a})}{R} - \frac{g}{4\pi} \frac{\mathbf{a} \times \hat{\mathbf{R}}}{R} \end{aligned} \quad (42)$$

in connection with the electromagnetic interaction, and also the Higgs scalar radiation described by

$$(D^i \phi)_{rad}^a = \frac{g_s}{4\pi} \frac{(\hat{\mathbf{R}} \cdot \mathbf{a}) \hat{\mathbf{R}}_i}{R} \hat{\mathbf{r}}'_a, \quad (D^0 \phi)_{rad}^a = \frac{g_s}{4\pi} \frac{\hat{\mathbf{R}} \cdot \mathbf{a}}{R} \hat{\mathbf{r}}'_a. \quad (43)$$

The duality is manifest in eq. (42); in fact, if one calculates the dyon radiation fields by invoking the duality argument to the well-known radiation fields of a uniformly accelerating electric charge, one ends up with eq. (42). Also the expressions in eq. (43) (more precisely, the quantities at front of $\hat{\mathbf{r}}'_a$) can be identified with the massless Klein-Gordon radiation fields produced by a uniformly accelerating point scalar source with strength $g_s = \pm \frac{4\pi}{e \cos \beta}$. Since the energy-momentum tensor is given by

$$T_{\mu\nu} = (D^\mu \phi)_a (D^\nu \phi)_a + G_a^{\mu\lambda} G_{a\lambda}^\nu + \eta^{\mu\nu} \mathcal{L}, \quad (44)$$

the radiation energy flux can be found using

$$\begin{aligned} T_{rad}^{0i} &= [G_a^{0k} G_a^{ik} + (D^0 \phi)^a (D^i \phi)^a]_{rad} \\ &= \mathbf{E}_{rad}^{em} \times \mathbf{B}_{rad}^{em} + (D^0 \phi)_{rad}^a (D^i \phi)_{rad}^a. \end{aligned} \quad (45)$$

Inserting eqs. (42) and (43) into this formula yields

$$T_{rad}^{0i} = \frac{(g^2 + q^2)}{16\pi^2 R^2} |\mathbf{a} \times \hat{\mathbf{R}}|^2 \hat{\mathbf{R}}_i + \frac{g_s^2}{16\pi^2 R^2} |\mathbf{a} \cdot \hat{\mathbf{R}}|^2 \hat{\mathbf{R}}_i, \quad (46)$$

where the first term is the electromagnetic contribution, and the second due to the Higgs scalar.

The dynamics of a BPS dyon in the presence of an asymptotically uniform electromagnetic field has been successfully extracted using only field equations of the Yang-Mills-Higgs system, at least in the small acceleration regime. The given field equations are intrinsically nonlinear, and they serve to specify the *entire* dynamics of a BPS dyon; the situation here is quite analogous to the case of Einstein's field equations [14]. The force law and radiation fields found by our analysis support the duality between electric and magnetic quantities. But note that quantum (or loop) effects have not been taken into account in our discussion.

Acknowledgements

Part of this work was done when one of us (C.L.) was visiting Center for Theoretical Physics, MIT during the summer of 1993, and he wishes to thank Prof. R. Jackiw for the hospitality especially and the support by the NSF on contract #910653. The work of D.B. is supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under contract #DE-AC02-76ER03069. The work of C.L. is supported in part by the Korean Science and Engineering Foundation (through the Center for Theoretical Physics, SNU) and the Ministry of Education, Republic of Korea.

References

- [1] G.'t Hooft, Nucl. Phys. B79 (1974) 276.
- [2] A.M. Polyakov, JETP Lett. 20 (1974) 194.
- [3] B. Julia and A. Zee, Phys. Rev. D11 (1975) 2227.
- [4] M.K. Prasad and C.M. Sommerfeld, Phys. Rev. Lett. 35 (1975) 760.
- [5] E.B. Bogomolny, Sov. J. Nucl. Phys. 24 (1976) 861.
- [6] S. Coleman, S. Parke, A. Neveu, and C.M. Sommerfeld, Phys. Rev. D15 (1977) 544.
- [7] N.S. Manton, Phys. Lett. B110 (1982) 54;
M.F. Atiyah and N.J. Hitchin, Phys. Lett. A107 (1985) 21;
G.W. Gibbons and N.S. Manton, Nucl. Phys. B274 (1986) 183.
- [8] D. Bak and C. Lee, Nucl. Phys. B403 (1993) 315.
- [9] N.S. Manton, Nucl. Phys. B126 (1977) 525.
- [10] N.S. Manton and T.M. Samols, Phys. Lett. B215 (1988) 559.
- [11] E. Weinberg, Phys. Rev. D20 (1979) 936.
- [12] R. Akhoury, J.-H. Jun and A.S. Goldhaber, Phys. Rev. D21 (1980) 454.
- [13] J.D. Jackson, Classical Electrodynamics, 2nd ed. (Wiley, New York, 1975), Chap.6.
- [14] A. Einstein, L. Infeld and B. Hoffman, Ann. Math. 39 (1938) 65.